arXiv:1202.2810v1 [math.MG] 13 Feb 2012

MANY NEIGHBORLY POLYTOPES AND ORIENTED MATROIDS

ARNAU PADROL

Abstract. We say that a d-polytope P is neighborly if every subset of $\lfloor \frac{d}{2} \rfloor$ vertices is a face of P. This concept applies naturally to oriented matroids. In 1982, Shemer introduced a sewing construction that allows to add a vertex to a neighborly polytope in such a way as to obtain a new neighborly polytope. We generalize this construction and extend it to oriented matroids.

Moreover we provide a new construction that allows to extend balanced oriented matroids. In the dual setting, this means that given a neighborly matroid of rank d with n elements, we obtain a new neighborly matroid of rank $d+1$ with $n+1$ elements. With the help of this new construction, we are able to generate many non-realizable neighborly matroids and to prove that the number of neighborly d-polytopes with n vertices is greater than $((n-1)/\mathrm{e}^{3/2})^{d(n-1)/2}$ when $n > 2d$.

1. Introduction

We say that a polytope is *k-neighborly* if every subset of vertices of size at most k is the set of vertices of one of its faces. It is easy to see that if a d-polytope is k-neighborly for any $k > \left\lfloor \frac{d}{2} \right\rfloor$, then it must be the d-dimensional simplex Δ_d. This is why we call a d-polytope *neighborly* if it is $\left\lfloor \frac{d}{2} \right\rfloor$-neighborly.

Neighborly polytopes are one of the most interesting families of polytopes because of their extremal properties. One of the main reasons for this is the Upper Bound Theorem by McMullen [20]: The number of i-dimensional faces of a d-polytope P with n vertices is maximal for neighborly polytopes, for all i.

The convex hull of any n points on the moment curve in $\mathbb{R}^d$, $\{(t, t^2, \ldots, t^d) : t \in \mathbb{R}\}$, determines a neighborly polytope: the cyclic polytope $c(n,d)$. The first non-cyclic neighborly polytope was found by Grünbaum in [15], who proved that for all $k \geq 2$ there are non-cyclic neighborly $(2k)$-polytopes with $2k+4$ vertices.

In 1981, Barnette [4] introduced the *facet splitting* technique, that allowed him to construct infinitely many neighborly polytopes, and to prove that the number of (combinatorial types of) neighborly d-polytopes with n vertices, $\mathrm{nb}(n,d)$, is bigger than

$$\mathrm{nb}(n,d) \geq \frac{(2n-4)!}{n!(n-2)!\binom{n}{d-3}} \sim 4^n.$$

This bound was improved by Shemer in [24], where he introduced the *sewing construction* to build an infinite family of neighborly polytopes in any even dimension. Given a neighborly d-polytope with n vertices and a suitable flag of faces, one can "sew" a new vertex onto it to

---

Date: February 14, 2012.

Supported by AGAUR grant 2009 SGR 1040, MEC grant MTM2008-03020, and FI grant from Catalunya's government DIUE and the ESF. Part of this work was done during a visit to ETH Zürich with support of Catalunya's government DIUE BE grant.

get a new neighborly d-polytope with $n+1$ vertices. With this construction, Shemer proved that $\mathrm{nb}(n,d)$ is greater than

$$\mathrm{nb}(n,d) \geq \frac{1}{2}\left(\left(\frac{d}{2}-1\right)\left\lfloor\frac{n-2}{d+1}\right\rfloor\right)! \sim 2^{c_d n \log n},$$

where $c_d \to \frac{1}{2}$ when $d \to \infty$.

This is a quite surprising result when it is combined with the fact that Goodman and Pollack proved in [14] that $\mathrm{nb}_l(n,d) \leq 2^{d(d+1)n\log n}$, where $\mathrm{nb}_l(n,d)$ is the number of (combinatorial types of) vertex-labeled d-polytopes with n vertices. Alon [1] improved this bound to

$$\mathrm{nb}_l(n,d) \leq \left(\frac{n}{d}\right)^{d^2 n(1+o(1))} \text{ when } \tfrac{n}{d} \to \infty.$$

Recently, Bisztriczky extended Shemer's technique to odd dimension [5], while Lee and Menzel provided a generalization of the sewing construction to non-simplicial polytopes [19]. In [26] details on how to compute the face lattice of a sewn polytope are given.

Oriented matroids are an abstract model for combinatorial geometry. In this sense, oriented matroids can be considered as a generalization of point configurations such as the vertex set of a convex polytope. The concept of neighborly polytope extends naturally to neighborly oriented matroids, see [10] or [25].

The first main result of this paper is Theorem 3.7, which is a generalization of the sewing construction to oriented matroids. It is a formulation valid for any rank (like in [5]) and extends to a bigger family of flags of faces that includes Barnette's facet splitting. Proposition 3.8 shows that in odd ranks, these are all the possible flags. The language of oriented matroids makes this proof of the sewing construction simpler than previous versions.

The major result in Section 4 is Theorem 4.2, which provides an alternative construction for neighborly matroids that works in the dual space. Given a neighborly matroid of rank d with n elements, we obtain a new neighborly matroid of rank $d+1$ with $n+1$ vertices. In a sense that we make precise in Section 5, this construction is a generalization of Shemer's.

Both techniques are based on the concept of *lexicographic extension of an oriented matroid*, which turns out to be a very useful tool for constructing polytopes with certain properties. Combining them, we provide non-realizable neighborly oriented matroids with n vertices and rank r for any $r \geq 5$ and $n \geq r+5$ in Theorem 5.4.

In the last section, Section 6, we use the dual construction to prove that the number of vertex-labeled neighborly polytopes with $r+d+1$ vertices and dimension d is greater than

$$\mathrm{nb}_l(r+d+1,d) \geq \frac{(r+d)^{\left(\frac{r}{2}+\frac{d}{2}\right)^2}}{r^{\left(\frac{r}{2}\right)^2} d^{\left(\frac{d}{2}\right)^2} \mathrm{e}^{3\frac{r}{2}\frac{d}{2}}}.$$

In particular, this bound is greater than $\left(\frac{n-1}{\mathrm{e}^{3/2}}\right)^{\frac{1}{2}d(n-1)}$ when $n > 2d$. This new bound not only improves Shemer's bound on the number of neighborly polytopes, but it is even greater than $\left(\frac{n-d}{d}\right)^{\frac{1}{4}nd}$, the current best lower bound on the number of all polytopes (valid only if $n \geq 2d$), found in [1].

2. Balanced and Neighborly Oriented Matroids

2.1. **Preliminaries.** We assume that the reader has some familiarity with the basics of oriented matroid theory; we refer to [12] for a nice introduction and to [6] for a comprehensive reference. We only present some results that we will directly use in our proofs.

As for notation, $\mathcal{M}$ will be an oriented matroid of rank d on a ground set E, with circuits $\mathcal{C}(\mathcal{M})$, cocircuits $\mathcal{C}^*(\mathcal{M})$, vectors $\mathcal{V}(\mathcal{M})$ and covectors $\mathcal{V}^*(\mathcal{M})$. Its dual $\mathcal{M}^*$ has rank $r = n-d$. $\mathcal{M}$ is uniform if the underlying matroid $\underline{\mathcal{M}}$ is uniform, that is, every subset of size d is a basis.

We view every vector (resp. covector) X indistinctly as a signed subset of E, $X = (X^+, X^-, X^0)$, or as a function from E to $\{+,-,0\}^n$. Hence, we will say $X(e) = +$ (or even $X(e) > 0$) meaning $e \in X^+$. $\underline{X}$ denotes the support $X^+ \cup X^-$ of X.

We say that two oriented matroids $\mathcal{M}_1$ and $\mathcal{M}_2$ on respective ground sets E_1 and E_2 are isomorphic, $\mathcal{M}_1 \simeq \mathcal{M}_2$, when there is a bijection between E_1 and E_2 that sends circuits of $\mathcal{M}_1$ to circuits of $\mathcal{M}_2$ (equivalently vectors, cocircuits or covectors).

A matroid $\mathcal{M}$ is *acyclic* if it contains $(+,+,\dots,+)$ as a covector. Its *facets* are the complements of the supports of its positive cocircuits, and its *faces* the complements of its positive covectors. Faces of rank 1 are called *vertices* of $\mathcal{M}$. In particular, every d-polytope is an acyclic matroid of rank $(d+1)$.

Similarly, a matroid is *totally cyclic* if it contains $(+,+,\dots,+)$ as a vector. We use the notation $\mathcal{D}_r$ for the only totally cyclic oriented matroid of rank r with $r+1$ elements.

Two elements $p, q \in E$ are *covariant* in $\mathcal{M}$ if they have the same sign in all circuits containing them; equivalently, if they have opposite signs in the cocircuits containing them. They are *contravariant* if they are covariant in the dual. We will say that p and q are $(+1)$-*inseparable* when they are covariant and that they are (-1)-*inseparable* when they are contravariant.

We will need some constructions to deal with an oriented matroid $\mathcal{M}$, in particular the *deletion* $\mathcal{M} \setminus e$ and the *contraction* $\mathcal{M}/e$ of an element e. They are defined by their covectors:

$$\mathcal{V}^*(\mathcal{M} \setminus e) = \left\{ V \in \{+,-,0\}^{n-1} \,\middle|\, (V,\sigma) \in \mathcal{V}^*(\mathcal{M}) \text{ for some } \sigma \in \{+,-,0\} \right\},$$
$$\mathcal{V}^*(\mathcal{M}/e) = \left\{ V \in \{+,-,0\}^{n-1} \,\middle|\, (V,0) \in \mathcal{V}^*(\mathcal{M}) \right\}.$$

Deletion and contraction are dual operations — $(\mathcal{M} \setminus e)^* = (\mathcal{M}^*/e)$ — that commute — $(\mathcal{M} \setminus p)/q = (\mathcal{M}/q) \setminus p$ — and naturally extend to subsets $S \subseteq E$ by iteratively deleting (resp. contracting) every element in S. An important observation is that $\mathcal{M} \setminus e$ and $\mathcal{M}/e$ determine $\mathcal{M}$ up to the reorientation of e:

Theorem 2.1 ([21, Theorem 4.1])**.** *Let $\mathcal{M}'$ and $\mathcal{M}''$ be two oriented matroids with $n-1$ elements, of respective ranks d and $d-1$, such that $\mathcal{V}^*(\mathcal{M}'') \subseteq \mathcal{V}^*(\mathcal{M}')$. Then there is an oriented matroid $\mathcal{M}$ with n elements with an special element p that fulfills $\mathcal{M} \setminus p = \mathcal{M}'$ and $\mathcal{M}/p = \mathcal{M}''$. The oriented matroid $\mathcal{M}$ has rank d and is unique up to reorientation of p.*

2.2. **Single Element Extensions.** Let $\mathcal{M}$ be an oriented matroid on a set E. A *single element extension* of $\mathcal{M}$ is an oriented matroid $\tilde{\mathcal{M}}$ on a ground set $\tilde{E} = E \cup p$ for some $p \notin E$, such that every circuit of $\mathcal{M}$ is a circuit in $\tilde{\mathcal{M}}$. Equivalently, $\tilde{\mathcal{M}}$ is a single element extension of $\mathcal{M}$ if $\mathcal{M}$ is a restriction of $\tilde{\mathcal{M}}$ by deleting one element. We will only consider extensions that do not increase the rank, *i.e.*, rank $(\tilde{\mathcal{M}}) = $ rank $(\mathcal{M})$.

Let $\tilde{\mathcal{M}} = \mathcal{M} \cup p$ be a single element extension of $\mathcal{M}$. Then, for every cocircuit $C = (C^+, C^-)$ of $\mathcal{M}$, there is a unique way to extend C to a cocircuit of $\tilde{\mathcal{M}}$: exactly one of $(C^+ \cup p, C^-)$,

$(C^+, C^- \cup p)$ or (C^+, C^-) is a cocircuit of $\tilde{\mathcal{M}}$. Hence, there is a unique function σ that assigns to each cocircuit C the value of $C(p) \in \{0, +, -\}$ in $\tilde{\mathcal{M}}$. We call such a function the *signature* of the extension.

Not every map from $\mathcal{C}^*(\mathcal{M})$ to $\{0, +, -\}$ corresponds to the signature of an extension. However, every valid signature uniquely determines the oriented matroid $\mathcal{M} \cup p$. In [6, Chapter 7] (originally from [17]) one can consult how to decide which signatures are valid, and how to determine all the cocircuits of $\tilde{\mathcal{M}}$ given the signature of the extension.

We will be interested in one particular family of single element extensions called lexicographic extensions.

Definition 2.2. Let $\mathcal{M}$ be an oriented matroid of rank r on a ground set E. Let $k \leq r$, let $(a_1, a_2, \dots, a_k)$ be an ordered subset of E and let $(s_1, s_2, \dots, s_k) \in \{+, -\}^k$ be a sign vector. The *lexicographic extension* of $\mathcal{M}$ by $p = [a_1^{s_1}, a_2^{s_2}, \dots, a_k^{s_k}]$ is defined as the single element extension that has the cocircuit signature

$$\sigma(C) = \begin{cases} s_i C(a_i) & \text{if } i \text{ is minimal with } C(a_i) \neq 0, \\ 0 & \text{if } C(a_i) = 0 \text{ for } i = 1, \dots, k. \end{cases}$$

We will often use the notation $\mathcal{M}[p]$ to denote a lexicographic extension $\mathcal{M} \cup p$.

Proposition 2.3. *If $\mathcal{M}$ is realizable, and $\mathcal{M}[p]$ is a lexicographic extension of $\mathcal{M}$, then $\mathcal{M}[p]$ is realizable.*

Proof. Let the vector configuration $V \subset \mathbb{R}^r$ be a realization of $\mathcal{M}$. If the signature of the lexicographic extension is $p = [v_1^{s_1}, v_2^{s_2}, \dots, v_k^{s_k}]$, let $v = s_1 v_1 + \varepsilon s_2 v_2 + \varepsilon^2 s_3 v_3 + \dots + \varepsilon^{k-1} s_k v_k$, for $\varepsilon > 0$ small enough. Then $V \cup v$ realizes $\mathcal{M} \cup p$. ☐

In the setting of a vector configuration V, the concept of lexicographic extension $p = [v_1^{s_1}, v_2^{s_2}, \dots, v_k^{s_k}]$ is very easy to understand. For every hyperplane h through vectors in $V \setminus \{v_1\}$, the new vector p must lie on the same side as $s_1 v_1$; for hyperplanes containing v_1 but not v_2, p must lie on the same side as $s_2 v_2$; *etc.*

Observe that if $\mathcal{M} \cup p$ is a lexicographic extension on $p = [a_1^{s_1}, \dots]$, then p and a_1 are always $(-s_1)$-inseparable.

Lexicographic extensions on uniform matroids behave well with respect to contractions. Proposition 2.6 can be seen as the restriction of [6, Proposition 7.1.2] to lexicographic extensions, and will be a very useful tool. Before proving it, we need a pair of results.

Lemma 2.4. *Let $\mathcal{M}$ be a uniform oriented matroid with two α-inseparable elements x and y.*

(1) For every circuit $X \in \mathcal{C}(\mathcal{M})$ with $X(x) = 0$ and $X(y) \neq 0$, there is a circuit $X' \in \mathcal{C}(\mathcal{M})$ with $X'(x) = -\alpha X(y)$, $X'(y) = 0$ and $X'(e) = X(e)$ for all $e \notin \{x, y\}$;

(2) For every cocircuit $C \in \mathcal{C}^*(\mathcal{M})$ with $C(x) = 0$ and $C(y) \neq 0$, there is a cocircuit $C' \in \mathcal{C}^*(\mathcal{M})$ with $C'(x) = \alpha C(y)$, $C'(y) = 0$ and $C'(e) = C(e)$ for all $e \notin \{x, y\}$.

Proof. Both statements are equivalent by duality. We prove the first one.

Let $X' \in \mathcal{C}(\mathcal{M})$ be the circuit with support $\underline{X'} = \underline{X} \setminus y \cup x$ and such that $X'(x) = -\alpha X(y)$. This circuit exists because $\mathcal{M}$ is uniform. We will see that $X'(e) = X(e)$ for all $e \in \underline{X'} \setminus x$. Let $e \in \underline{X} \setminus y$, and let C be the cocircuit of $\mathcal{M}$ with $\underline{C} = E \setminus (\underline{X} \setminus y \setminus e)$. That makes $\underline{C} \cap \underline{X} = \{e, y\}$. Since y and x are α-inseparable, $C(x) = -\alpha C(y)$, and by circuit-cocircuit orthogonality,

$$X(y)X(e) = -C(y)C(e) = \alpha C(x)C(e).$$

But $\underline{C} \cap \underline{X}' = \{e, x\}$, and hence, again by orthogonality, $X'(x)X'(e) = -C(x)C(e)$. The conclusion now follows from $X'(x) = -\alpha X(y)$. □

Lemma 2.5. *If $\mathcal{M}$ is uniform and $p = [a_1^{s_1}, \dots]$, then*

$$\mathcal{M}/a_1 = (\mathcal{M}[p] \setminus p)/a_1 = (\mathcal{M}[p] \setminus a_1)/p.$$

Proof. The first equality is direct. The second one follows from Lemma 2.4. Indeed, every cocircuit of $(\mathcal{M}[p]/a_1) \setminus p$ corresponds to a cocircuit C of $\mathcal{M}[p]$ with $C(a_1) = 0$ and $C(p) \neq 0$. By Lemma 2.4, the values of C on $e \notin \{a_1, p\}$ coincide with the values of C' on $e \notin \{a_1, p\}$, where C' is a cocircuit of $\mathcal{M}[p]$ with $C'(a_1) \neq 0$ and $C'(p) = 0$. That is, C' corresponds to a cocircuit of $(\mathcal{M}[p]/p) \setminus a_1$. □

Proposition 2.6. *Let $\mathcal{M}$ be a uniform oriented matroid of rank d on a ground set E, and $\tilde{\mathcal{M}} = \mathcal{M} \cup p$ a lexicographic extension by $p = [a_1^{s_1}, a_2^{s_2}, \dots, a_d^{s_d}]$. Then*

$$\tilde{\mathcal{M}}/p \simeq (\mathcal{M}/a_1)[a_2^{-s_1 s_2}, \dots, a_d^{-s_1 s_d}], \tag{1}$$

$$\tilde{\mathcal{M}}/a_i = (\mathcal{M}/a_i)[a_1^{s_1}, \dots, a_{i-1}^{s_{i-1}}, a_{i+1}^{s_{i+1}}, \dots, a_d^{s_d}], \tag{2}$$

$$\tilde{\mathcal{M}}/e = (\mathcal{M}/e)[a_1^{s_1}, a_2^{s_2}, \dots, a_{d-1}^{s_{d-1}}], \tag{3}$$

where $e \in E$ is any element different from p and any a_i, and the isomorphism φ in eq. (1) *is $\varphi(a) = a$ for all $a \in E \setminus \{p, a_1\}$ and $\varphi(a_1) = [a_2^{-s_1 s_2}, \dots, a_d^{-s_1 s_d}]$ is the extending element.*

Proof. The proof of (2) and (3) is direct just by observing the signature of p in $\tilde{\mathcal{M}}$.

To prove the case $\tilde{\mathcal{M}}/p$, observe that $(\tilde{\mathcal{M}}/p) \setminus a_1 = \mathcal{M}/a_1$ by Lemma 2.5. Therefore, we only need to prove that the signature of a_1 in $(\tilde{\mathcal{M}}/p) \setminus a_1$ coincides with that of the lexicographic extension $[a_2^{-s_1 s_2}, \dots, a_d^{-s_1 s_d}]$. That is, let $C \in \mathcal{C}^*(\tilde{\mathcal{M}})$ be a cocircuit of $\tilde{\mathcal{M}}$ with $C(p) = 0$ and $C(a_1) \neq 0$ and let $k > 1$ be minimal with $C(a_k) \neq 0$. We want to see that $C(a_1) = -s_1 s_k C(a_k)$.

Because a_1 and p are $(-s_1)$-inseparable, Lemma 2.4 yields a cocircuit $C' \in \mathcal{C}^*(\tilde{\mathcal{M}})$ with $C'(p) = -s_1 C(a_1)$ and $C'(a_1) = 0$ and such that k is minimal with $C'(a_k) \neq 0$. Moreover $C'(a_k) = C(a_k)$ and by the signature of the lexicographic extension $C'(p) = s_k C'(a_k) = s_k C(a_k)$. The claim follows from comparing both expressions for $C'(p)$. □

Since (2) and (3) are direct, the most interesting case of this lemma is the first one, namely $\tilde{\mathcal{M}}/p \simeq (\mathcal{M}/a_1)[a_2^{-s_1 s_2}, \dots, a_d^{-s_1 s_d}]$. The idea behind it is that every hyperplane that goes through p and not through a_1 looks very much like some hyperplane that goes through a_1 and not p. If $s_1 = +$, then a_1 and p are very close, which means that when we perturb a hyperplane h that goes through a_1 with p in h^+ to its analog h' through p, then a_1 lies in h'^-. On the other hand, if $s_1 = -$, then a_1 and $-p$ are very close, and to perturb h to h', one must also switch its orientation. Hence if p was in h^+, then a_1 is in h'^-.

2.3. **Balanced and neighborly oriented matroids.**

The definition of neighborliness can be generalized to oriented matroids, so that neighborly polytopes correspond to realizable neighborly oriented matroids.

Definition 2.7. Let $\mathcal{M}$ be an oriented matroid of rank d on a ground set E. We say that $\mathcal{M}$ is *neighborly* if every subset $R \subset E$ of size at most $\left\lfloor \frac{d-1}{2} \right\rfloor$ is a face of $\mathcal{M}$. That is, there must exist a covector C of $\mathcal{M}$ with $C^+ = E \setminus R$ and $C^- = \emptyset$.

In particular, every neighborly matroid is acyclic and all of its elements are vertices. An interesting property of neighborly matroids of odd rank (hence polytopes of even dimension) is that they are rigid. We call an oriented matroid $\mathcal{M}$ *rigid* if there is no other matroid that shares the same face lattice with $\mathcal{M}$.

Theorem 2.8 ([25, Theorem 4.2])**.** *Every neighborly oriented matroid of rank* $2m+1$ *is rigid.*

Another important concept for this paper is that of balanced matroid.

Definition 2.9. An oriented matroid of rank d $\mathcal{M}$ is *balanced* if every cocircuit C of $\mathcal{M}$ is *halving*, *i.e.*,

$$\left\lfloor \frac{n-d+1}{2} \right\rfloor \leq |C^+| \leq \left\lceil \frac{n-d+1}{2} \right\rceil .$$

Let $\mathcal{M}$ be an oriented matroid. The *discrepancy* of a cocircuit C of $\mathcal{M}$ is $\operatorname{disc}(C) = ||C^+| - |C^-||$, and the discrepancy of $\mathcal{M}$ is the maximal discrepancy among its cocircuits, $\operatorname{disc}(\mathcal{M}) = \max_{C\in\mathcal{C}^*(\mathcal{M})} \operatorname{disc}(C)$. In a uniform oriented matroid, a cocircuit C is halving if and only if it has discrepancy 0 or 1, and a uniform oriented matroid $\mathcal{M}$ is balanced if and only if it has discrepancy 0 or 1.

A key result is that neighborliness and balancedness are dual concepts:

Proposition 2.10 ([25, Proposition 3.2])**.** *An oriented matroid* $\mathcal{M}$ *is neighborly if and only if its dual matroid* $\mathcal{M}^*$ *is balanced.*

We will also use the following lemma.

Lemma 2.11. *If* $\mathcal{M}$ *is a uniform balanced oriented matroid of rank* $r \geq 2$ *with* n *elements such that* $n-r-1$ *is even, then all inseparable elements in* $\mathcal{M}$ *must be covariant.*

Analogously, if $\mathcal{P}$ *is a uniform neighborly oriented matroid of odd rank* $2m+1$ *with at least* $2m+3$ *elements, then all inseparable elements in* $\mathcal{P}$ *must be contravariant.*

Proof. Both results are equivalent by duality. To prove the second claim, observe that if p and q are α-inseparable in $\mathcal{P}$, then they are also in $\mathcal{P} \setminus S$ for any S that contains neither p nor q. Hence we can remove elements from $\mathcal{P}$ until we are left with a neighborly matroid with $2m+3$ elements. All neighborly matroids of even dimension and corank 2 are cyclic polytopes, and those only have contravariant pairs. □

3. The sewing construction

This section is devoted to explaining the sewing construction, introduced by Shemer in [24], that allows to construct an infinite class of neighborly polytopes. Even if Shemer described the sewing construction in terms of Grünbaum's *beneath-beyond* technique, it is in fact a lexicographic extension, and we will explain it in these terms. In this section, we use the letter $\mathcal{P}$ for oriented matroids to reinforce the idea that all the following results translate directly to polytopes.

Let $\mathcal{P}$ be an acyclic oriented matroid and let F be a facet of $\mathcal{P}$. That is, there exists a cocircuit C_F of $\mathcal{P}$ such that $C_F(e) = 0$ if $e \in F$ and $C_F(e) = +$ otherwise. Let p be a single element extension of $\mathcal{P}$ with signature σ_p. We say that p is *beneath* F if $\sigma_p(C_F) = +$, that p is *beyond* F when $\sigma_p(C_F) = -$, and that p is *on* F if $\sigma_p(C_F) = 0$. We say that an extension p of $\mathcal{P}$ lies *exactly beyond* a set of facets $\mathcal{F}$ if it lies beyond all facets in $\mathcal{F}$ and beneath all facets not in $\mathcal{F}$.

It is enough to know the signature of p on the facet cocircuits of $\mathcal{P}$ to tell the whole face lattice of $\mathcal{P} \cup p$.

A *flag* of an acyclic oriented matroid $\mathcal{P}$ is a strictly increasing sequence of proper faces $T_1 \subset T_2 \subset \cdots \subset T_k$. Given a flag $\mathcal{T} = \{T_j\}_{j=1}^k$ of $\mathcal{P}$, let $\mathcal{F}_j$ be the set of facets of $\mathcal{P}$ that contain T_j, and let $\mathcal{F}(\mathcal{T}) := \mathcal{F}_1 \setminus (\mathcal{F}_2 \setminus (\cdots \setminus \mathcal{F}_k) \dots)$; that is

$$\mathcal{F}(\mathcal{T}) = \begin{cases} (\mathcal{F}_1 \setminus \mathcal{F}_2) \cup (\mathcal{F}_3 \setminus \mathcal{F}_4) \cup \cdots \cup (\mathcal{F}_{k-1} \setminus \mathcal{F}_k) & \text{if } k \text{ is even,} \\ (\mathcal{F}_1 \setminus \mathcal{F}_2) \cup (\mathcal{F}_3 \setminus \mathcal{F}_4) \cup \cdots \cup \mathcal{F}_k & \text{if } k \text{ is odd.} \end{cases}$$

Given a polytope P with a flag $\mathcal{T} = T_1 \subset T_2 \subset \cdots \subset T_k$, Shemer proved that there always exists an extension exactly beyond $\mathcal{F}(\mathcal{T})$, via the sewing construction. We will show that there is a lexicographic extension that realizes the desired signature.

Definition 3.1 (Sewing onto a flag)**.** Let $\mathcal{T} = \{T_j\}_{j=1}^k$ be a flag of an acyclic matroid $\mathcal{P}$. We extend it with $T_{k+1} = \mathcal{P}$ and define $U_i = T_i \setminus T_{i-1}$. We say that p is *sewn* onto $\mathcal{P}$ through $\mathcal{T}$, if p is a lexicographic extension

$$p = [T_1^+, U_2^-, U_3^+, \dots, U_{k+1}^{(-1)^k}],$$

where these sets represent their elements in any order.

That is, p is a lexicographic extension $p = [a_1^{s_1}, a_2^{s_2}, \dots, a_n^{s_n}]$ where $a_1, \dots, a_n$ are the points in T_{k+1}, sorted such that if a_i belongs to a T_s and a_j does not then $i < j$, and where s_j is $+$ if the smallest i such that $x_j \in T_i$ is odd, and $-$ otherwise.

We use the notation $\mathcal{P}[\mathcal{T}]$ to designate the extension $\mathcal{P} \cup p$ when p is sewn onto $\mathcal{P}$ through $\mathcal{T}$.

In terms of oriented matroids, the definition of $\mathcal{P}[\mathcal{T}]$ is ambiguous, since it can represent different oriented matroids. However, the following proposition shows that all the extensions $\mathcal{P}[\mathcal{T}]$ have the same face lattice. Moreover, because of rigidity, this implies that there is no ambiguity when $\mathcal{P}[\mathcal{T}]$ is neighborly of odd rank.

Proposition 3.2. *Let $\mathcal{T} = \{T_j\}_{j=1}^k$ be a flag of an acyclic oriented matroid $\mathcal{P}$, and let $\mathcal{F}(\mathcal{T}) := \mathcal{F}_1 \setminus (\mathcal{F}_2 \setminus (\cdots \setminus \mathcal{F}_k) \dots)$. If p is the lexicographic extension $\mathcal{P} \cup p = \mathcal{P}[\mathcal{T}]$, then p lies exactly beyond $\mathcal{F}$.*

Proof. Let p be the lexicographic extension $[a_1^{s_1}, a_2^{s_2}, \dots, a_{d+1}^{s_{d+1}}]$ from Definition 3.1. We have to see that p lies beneath any facet in $\mathcal{F}_{2j} \setminus \mathcal{F}_{2j+1}$ and beyond any facet in $\mathcal{F}_{2j-1} \setminus \mathcal{F}_{2j}$.

If C is a cocircuit representing a facet F in $\mathcal{F}_j \setminus \mathcal{F}_{j+1}$, then the first a_i with $C(a_i) \neq 0$ belongs to T_{j+1}. If j is even then $s_i = +$ and hence p is beneath F. If j is odd then $s_i = -$ and p is beyond F. □

The sewing construction starts with a neighborly oriented matroid $\mathcal{P}$ of rank d with n elements and gives a neighborly oriented matroid $\tilde{\mathcal{P}}$ of rank d with $n+1$ elements, provided that $\mathcal{P}$ has a *universal flag*.

Definition 3.3. Let $\mathcal{P}$ be a uniform acyclic oriented matroid on E of rank d, and let $m = \left\lfloor \frac{d-1}{2} \right\rfloor$.

(i) A face F of $\mathcal{P}$ is a *universal face* if the contraction $\mathcal{P}/F$ is neighborly.
(ii) A flag $\mathcal{T}$ of $\mathcal{P}$ is a *universal flag* if $\mathcal{T} = \{T_j\}_{j=1}^m$ where each T_j is a universal face with $2j$ vertices.

The first example of neighborly polytopes with universal flags are cyclic polytopes (see [24, Theorem 3.4]).

Example 3.4. Let $c(n, 2m)$ be a cyclic polytope of even dimension $2m$, with vertices $s_1, \dots, s_n$ labeled in cyclic order. Then $\{s_i, s_{i+1}\}$, $1 \leq i < n$ and $\{s_1, s_n\}$ are universal edges of $c(n, 2m)$. Moreover, $c(n, 2m)/\{s_i, s_{i+1}\} \simeq c(n-2, 2m-2)$ with the same cyclic order. In particular this gives a large family of universal flags of $c(n, 2m)$ formed by faces that are the union of a universal edge of $c(n, 2m)$ with a (possibly empty) universal face of $c(n-2, 2m-2)$.

With these notions, we are ready to present Shemer's Sewing Theorem [24, Theorem 4.6].

Theorem 3.5 (The Sewing Theorem). *Let P be a neighborly polytope of dimension $2m$ with a universal flag $\mathcal{T} = \{T_j\}_{j=1}^m$, where $T_j = \{x_i, y_i\}_{i=1}^j$. Let $\tilde{P} = P[\mathcal{T}]$, with p sewn onto P through $\mathcal{T}$. Then,*

1. *$\tilde{P}$ is a neighborly polytope with vertices* $\operatorname{vert}(\tilde{P}) = \operatorname{vert}(P) \cup \{p\}$.
2. *For all $1 \leq j \leq m$, $T_{j-1} \cup \{x_j, p\}$ and $T_{j-1} \cup \{y_j, p\}$ are universal faces of $\tilde{P}$. If moreover j is even, then T_j is also a universal face of $\tilde{P}$.*

Combining Example 3.4 and Theorem 3.5, one can obtain a large family of neighborly polytopes: starting with an even-dimensional cyclic polytope and one of its universal flags, sew a new vertex onto it and obtain a new neighborly polytope (point 1 of Theorem 3.5) with new universal flags (from point 2 of Theorem 3.5). Since the new polytope has universal flags, one can keep sewing again. This method generates a family of neighborly polytopes $\mathcal{S}$, that we call *totally sewn* polytopes[1].

In Theorem 3.7 we generalize Theorem 3.5 to neighborly oriented matroids. Our formulation, like [5, Theorem 2], does not depend on the parity of the rank. Moreover, it is extended to any flag that contains a universal subflag, as suggested in [24, Remark 7.4]. In order to prove it, we need a lemma.

We will use the notation $\mathcal{T}'/T_i = \{T'_j/T_i\}_{j=i+1}^m$ where T'_j/T_i is the face of $\mathcal{P}/T_i$ that represents T'_j.

Lemma 3.6. *Let $\mathcal{P}$ be a uniform neighborly matroid of rank d. Let $\mathcal{T}' = \{T'_j\}_{j=1}^k$ be a flag of $\mathcal{P}$ that contains a universal subflag $\mathcal{T} = \{T_j\}_{j=1}^m$, where $m = \left\lfloor \frac{d-1}{2} \right\rfloor$ and $T_j = \{x_i, y_i\}_{i=1}^j$. Let p be sewn onto $\mathcal{P}$ through $\mathcal{T}'$. If $T_{i-1} \cup \{y_i\}$ does not belong to $\mathcal{T}'$, then*

$$\mathcal{P}[\mathcal{T}']/\{T_{i-1}, x_i, p\} \simeq (\mathcal{P}/T_i)[\mathcal{T}'/T_i].$$

The isomorphism sends y_i to the sewn vertex and the remaining are the natural mappings.

Proof. By Proposition 2.6, the contraction $\mathcal{P}[\mathcal{T}']/T_{i-1}$ is a lexicographic extension of $\mathcal{P}/T_{i-1}$ whose signature coincides with that of $[\mathcal{T}']$ by removing the first $2(i-1)$ elements. Hence $\mathcal{P}[\mathcal{T}']/T_{i-1}$ must be one of the extensions

$$\mathcal{P}[\mathcal{T}']/T_{i-1} = \begin{cases} \mathcal{P}/T_{i-1}[x_i^+, y_i^+, x_{i+1}^-, \dots], \\ \mathcal{P}/T_{i-1}[x_i^-, y_i^-, x_{i+1}^+, \dots], \\ \mathcal{P}/T_{i-1}[x_i^+, y_i^-, x_{i+1}^+, \dots], \\ \mathcal{P}/T_{i-1}[x_i^-, y_i^+, x_{i+1}^-, \dots]. \end{cases}$$

If T'_{j-1} is the face of $\mathcal{T}'$ corresponding to T_{i-1}, and $U'_j = T'_j \setminus T'_{j-1}$, then the first two cases are possible when $U'_j = \{x_i, y_i\}$ and the last two cases are possible when $U'_j = \{x_i\}$.

---

[1] Shemer's concept of totally sewn polytopes is slightly different, because we only admit those flags that come from Theorem 3.5.

We use Proposition 2.6 twice on each of these (contracting successively x_i and p) to get $\mathcal{P}[\mathcal{T}']/\{T_{i-1}, x_i, p\} \simeq (\mathcal{P}/\{T_{i-1}, x_i, y_i\})[x_{i+1}^+, \dots] = (\mathcal{P}/T_i)[\mathcal{T}'/T_i]$. □

We can finally state and prove the Extended Sewing Theorem.

Theorem 3.7 (The Extended Sewing Theorem)**.** *Let $\mathcal{P}$ be a uniform neighborly oriented matroid of rank d with a universal flag $\mathcal{T} = \{T_j\}_{j=1}^m$, where $T_j = \{x_i, y_i\}_{i=1}^j$ and $m = \lfloor \frac{d-1}{2} \rfloor$. Let $\mathcal{T}' = \{T_j'\}_{j=1}^k$ be a flag of $\mathcal{P}$ that contains $\mathcal{T}$ as a subflag. Let $\tilde{\mathcal{P}} = \mathcal{P}[\mathcal{T}']$, with p sewn onto $\mathcal{P}$ through $\mathcal{T}'$. Then $\tilde{\mathcal{P}}$ is a uniform neighborly matroid of rank d.*

Proof. Depending on whether $T_1' = \{x_1, y_1\}$ or $T_1' = \{x_1\}$, the extension $\tilde{\mathcal{P}} = \mathcal{P} \cup p$ is either the lexicographic extension $\tilde{\mathcal{P}} = \mathcal{P}\left[x_1^+, y_1^+, \dots\right]$, or $\tilde{\mathcal{P}} = \mathcal{P}\left[x_1^+, y_1^-, \dots\right]$.

We check that $\tilde{\mathcal{P}}$ is neighborly by checking that $\tilde{\mathcal{P}}^*$ is balanced, *i.e.*, we check that every circuit X of $\tilde{\mathcal{P}}$ is halving:

(1) If $X(p) = 0$, then X is halving, since it is a circuit of $\mathcal{P}$, and $\mathcal{P}$ is neighborly.
(2) If $X(p) \neq 0$ and $X(x_1) = 0$, we use that p and x_1 are contravariant. By Lemma 2.4, if $X' \in \mathcal{C}(\tilde{\mathcal{P}})$ is the circuit with $X'(x_1) = X(p)$ and support $\underline{X'} = \underline{X} \setminus p \cup x_1$, then $|X^+| = |X'^+|$ and $|X^-| = |X'^-|$. Since $X'(p) = 0$, X' is halving by the previous point. Hence so is X.
(3) If $X(p) \neq 0$ and $X(x_1) \neq 0$, since p and x_1 are contravariant, $X(p) = -X(x_1)$. Observe that the rest of the values of X correspond to a circuit of $\tilde{\mathcal{P}}/\{p, x_1\}$. If $\tilde{\mathcal{P}}/\{p, x_1\}$ is balanced, we are done.

    By Lemma 3.6, $\tilde{\mathcal{P}}/\{p, x_1\} \simeq \mathcal{P}/T_1[\mathcal{T}'/T_1]$. Since $\{x_1, y_1\}$ was a universal edge, $\mathcal{P}/T_1$ is a neighborly matroid and $\mathcal{T}'/T_1$ is a flag that contains the universal flag $\mathcal{T}/T_1$. Then the result follows by induction on d. Observe for the base case that all acyclic universal matroids of rank 1 or 2 are neighborly.

□

For matroids of odd rank, the flags of the previous theorem are the only ones that can yield a neighborly polytope.

Proposition 3.8. *Let $\mathcal{P}$ be a uniform neighborly oriented matroid of rank $d = 2m + 1$ with more than $d+1$ elements and with a flag $\mathcal{T}$, and let $\tilde{\mathcal{P}} = \mathcal{P}[\mathcal{T}]$, with p sewn onto $\mathcal{P}$ through $\mathcal{T}$. Then $\tilde{\mathcal{P}}$ is neighborly if and only if $\mathcal{T}$ contains a universal subflag.*

Proof. By Theorem 3.7, this condition is sufficient. To find necessary conditions, we use that $\tilde{\mathcal{P}}$ is neighborly if and only if every circuit of $\tilde{\mathcal{P}}$ is halving.

By definition, $\tilde{\mathcal{P}} = \mathcal{P}[\mathcal{T}] = \mathcal{P} \cup p$ is a lexicographic extension that has the form $\tilde{\mathcal{P}} = \mathcal{P}\left[a_1^+, a_2^{s_2}, \dots, a_d^{s_d}\right]$. Let $X \in \mathcal{C}(\tilde{\mathcal{P}})$ be a circuit with $\{p, a_1\} \subset \underline{X}$. Since p and a_1 are contravariant, $X(p) = -X(a_1)$. Hence, if X is balanced, so must be $X \setminus \{p, a_1\}$. Observe that $X \setminus \{p, a_1\}$ is a circuit of $\tilde{\mathcal{P}}/\{p, a_1\}$ and that all the circuits of $\tilde{\mathcal{P}}/\{p, a_1\}$ arise this way. Hence $\tilde{\mathcal{P}}/\{p, a_1\}$ must be neighborly.

By Proposition 2.6,

$$\tilde{\mathcal{P}}/\{p, a_1\} \simeq \mathcal{P}/\{a_1, a_2\}\left[a_3^{-s_2 s_3},\ a_4^{-s_2 s_4},\ , \dots,\ a_d^{-s_2 s_d}\right].$$

Observe that a_2 and a_3 are $(s_2 s_3)$-inseparable in $\tilde{\mathcal{P}}/\{p, a_1\}$, which is a neighborly matroid of odd rank and corank at least 2. By Lemma 2.11, $s_2 s_3 = -$.

In particular, either $s_2 = +$ and $s_3 = -$, or $s_2 = -$ and $s_3 = +$. The first option implies that $T_1 = \{a_1, a_2\}$, while the second option implies that $T_1 = \{a_1\}$ and $T_2 = \{a_1, a_2\}$. Moreover, if $\tilde{\mathcal{P}}/\{p, a_1\}$ is neighborly, then $(\tilde{\mathcal{P}}/\{p, a_1\}) \setminus a_2 \simeq \mathcal{P}/\{a_1, a_2\}$ must be neighborly and hence $\{a_1, a_2\}$ must be a universal edge of $\mathcal{P}$.

Now the result follows by induction on d. For the base case just observe that neighborly matroids of rank 3 are polygons, and the only flags that yield a polygon with one extra vertex are of the form $\{x\} \subset \{x, y\}$ or just $\{x, y\}$, where $\{x, y\}$ is an edge of the polygon. $\square$



FIGURE 1. Sewing onto $\{A, E\}$ (middle) and sewing onto $\{A\} \subset \{A, E\}$ (right). In the first case, $\{A, F\}$ and $\{E, F\}$ become universal faces, and $\{A, E\}$ is not a universal face any more. In the second case, $\{A, F\}$ and $\{A, E\}$ are universal faces, while $\{E, F\}$ is not.

We can tell many universal faces (and flags) of the neighborly polytopes constructed using Theorem 3.7 thanks to Proposition 3.9, which is the analogue of point 2 of Theorem 3.5.

We use the following notation. If $\mathcal{T}' = \{T'_j\}_{j=1}^{k}$ is a flag of $\mathcal{P}$ that contains a universal subflag $\mathcal{T} = \{T_j\}_{j=1}^{m}$, where $T_j = \{x_i, y_i\}_{i=1}^{j}$, we will say that T_i is *x_i-split* (resp. *y_i-split*) if $T_{i-1} \cup \{x_i\}$ (resp. $T_{i-1} \cup \{y_i\}$) belongs to $\mathcal{T}'$.

Moreover, we say that T_i is *even* if the number of non-split faces T_j with $j \le i$ is even, otherwise we say that T_i is *odd*.

Proposition 3.9. *Let $\mathcal{P}$ be a uniform neighborly oriented matroid of rank d with a flag $\mathcal{T}' = \{T'_j\}_{j=1}^{k}$ that contains a universal subflag $\mathcal{T} = \{T_j\}_{j=1}^{m}$, where $T_j = \{x_i, y_i\}_{i=1}^{j}$ and $m = \left\lfloor \frac{d-1}{2} \right\rfloor$. Let $\tilde{\mathcal{P}} = \mathcal{P}[\mathcal{T}']$, with p sewn onto $\mathcal{P}$ through $\mathcal{T}'$. Then the following are universal faces of $\tilde{\mathcal{P}}$:*

(1) *T_i, where $1 \le i \le m$, if T_i is even.*
(2) *$(T_j \setminus x_i) \cup p$, where $1 \le i \le j \le m$, if*
  (i) *T_i is not split and T_j/T_i is even, or*
  (ii) *T_i is x_i-split and T_j/T_i is odd, or*
  (iii) *T_i is y_i-split and T_j/T_i is even.*
(3) *$(T_j \setminus y_i) \cup p$, where $1 \le i \le j \le m$, if*
  (i) *T_i is not split and T_j/T_i is even, or*
  (ii) *T_i is x_i-split and T_j/T_i is even, or*
  (iii) *T_i is y_i-split and T_j/T_i is odd.*

Proof. Without loss of generality, we will assume that all split faces are x_i-split. The proof relies on applying, case by case, Proposition 2.6 to reduce the contraction to a lexicographic extension that we know to be neighborly because of Theorem 3.7. We also use the following observation about the signature of the extension $\mathcal{P}[\mathcal{T}']$: the sign of x_i is $+$ if and only if T_{i-1} is even. This follows from the fact that

$$\tilde{\mathcal{P}} = \begin{cases} \mathcal{P}[\dots, a^{\alpha}, x_i^{-\alpha}, y_i^{\alpha}, b^{-\alpha}, \dots] & \text{if } T_i \text{ is } x_i\text{-split,} \\ \mathcal{P}[\dots, a^{\alpha}, x_i^{-\alpha}, y_i^{-\alpha}, b^{\alpha}, \dots] & \text{if it is not split.} \end{cases}$$

In particular, if T_i is even, $\tilde{\mathcal{P}}/T_i = \mathcal{P}/T_i[\mathcal{T}'/T_i]$, and T_i is a universal flag. This proves point 1. Moreover, independently of whether T_i is even or odd,

$$\tilde{\mathcal{P}}/(T_{i-1} \cup \{p\}) \simeq \begin{cases} \mathcal{P}/(T_{i-1} \cup \{x_i\})[y_i^{+}, x_{i+1}^{-}, \dots] & \text{if } T_i \text{ is split,} \\ \mathcal{P}/(T_{i-1} \cup \{x_i\})[y_i^{-}, x_{i+1}^{+}, \dots] & \text{if it is not.} \end{cases}$$

Hence, $\tilde{\mathcal{P}}/(T_{i-1} \cup \{x_i, p\}) \simeq \mathcal{P}/T_i[\mathcal{T}'/T_i]$. If T_i is not split $\tilde{\mathcal{P}}/(T_{i-1} \cup \{y_i, p\}) \simeq \mathcal{P}/T_i[\mathcal{T}'/T_i]$. Moreover, we know that T_j/T_i is a universal face of $\mathcal{P}/T_i[\mathcal{T}'/T_i]$ when it is even. This proves points 2i, 2iii, 3i and 3ii.

If T_i is split, then $\tilde{\mathcal{P}}/(T_{i-1} \cup \{y_i, p\}) \simeq \mathcal{P}/T_i[-\mathcal{T}'/T_i]$, where $[-\mathcal{T}'/T_i]$ means the extension $[\mathcal{T}'/T_i]$ with the signs reversed. And using the previous observation, $\mathcal{P}/T_i[-\mathcal{T}'/T_i]/(T_j/T_i) \simeq \mathcal{P}/T_j[\mathcal{T}'/T_j]$ when T_j/T_i is odd. Which proves the remaining points 2ii and 3iii. ☐

In particular, Proposition 3.9 provides a simple way to tell universal flags of $\mathcal{P}[\mathcal{T}']$. If T_1 is not split then $\{x_1, p\}$ and $\{y_1, p\}$ are universal edges of $\mathcal{P}[\mathcal{T}']$; if it is x_1-split, then $\{x_1, p\}$ and $\{x_1, y_1\}$ are universal edges of $\mathcal{P}[\mathcal{T}']$; and if it is y_1-split, then $\{y_1, p\}$ and $\{x_1, y_1\}$ are universal edges of $\mathcal{P}[\mathcal{T}']$. Contracting any of these edges leaves a matroid isomorphic to $\mathcal{P}/T_1[\mathcal{T}'/T_1]$, and we can inductively build a universal flag of $\mathcal{P}[\mathcal{T}']$. The rank 3 example in Figure 1 can give some intuition on why these universal edges appear.

Example 3.10. Let $\mathcal{M}$ be a neighborly oriented matroid of rank 5 with a universal tower $\mathcal{T} = \{a, b\} \subset \{a, b, c, d\}$. Consider the following lexicographic extensions

$$\begin{aligned} p_1 &= [a^+, b^+, c^-, d^-, e^+], \\ p_2 &= [a^+, b^-, c^+, d^+, e^-], \\ p_3 &= [a^+, b^+, c^-, d^+, e^-] \text{ and} \\ p_4 &= [a^+, b^-, c^+, d^-, e^+], \end{aligned}$$

where e is any element of $\mathcal{M}$. They give rise to the oriented matroids $\mathcal{M}_1 = \mathcal{M}[p_1]$, $\mathcal{M}_2 = \mathcal{M}[p_2]$, $\mathcal{M}_3 = \mathcal{M}[p_3]$ and $\mathcal{M}_4 = \mathcal{M}[p_4]$.

The following table shows for which $\mathcal{M}_i$ is each of the following sets of vertices a universal face.

	ab	p_ib	ap_i	$abcd$	p_ibcd	ap_icd	abp_id	$abcp_i$
$\mathcal{M}_1$	✗	✓	✓	✓	✗	✗	✓	✓
$\mathcal{M}_2$	✓	✗	✓	✗	✓	✗	✓	✓
$\mathcal{M}_3$	✗	✓	✓	✗	✓	✓	✗	✓
$\mathcal{M}_4$	✓	✗	✓	✓	✗	✓	✗	✓

Remark 3.11. Theorem 3.7 not only generalizes the Sewing Theorem. It also includes Barnette's facet-splitting technique [4, Theorem 3].

Just like we did to construct the family $\mathcal{S}$, we can combine Example 3.4, Theorem 3.7 and Proposition 3.9 to obtain a large family of neighborly polytopes that contains $\mathcal{S}$. We call this family $\mathcal{E}$. In fact, since [19, Theorem 5.1] implies that cyclic polytopes belong to $\mathcal{E}$, it is enough to start sewing on a simplex.

Moreover, since every subpolytope of a neighborly polytope is a neighborly polytope, any polytope obtained from a polytope in $\mathcal{E}$ and then omitting some vertices is also a neighborly polytope. The polytopes that can be obtained this way via sewing and omitting form a family that we denote $\mathcal{O}$.

4. THE GALE SEWING CONSTRUCTION

We present here a different construction, also based on lexicographic extensions, that allows us to construct neighborly matroids. This construction works in the dual, that is, it extends balanced matroids to new balanced matroids.

Proposition 4.1. *Let $\mathcal{M}$ be a uniform oriented matroid of rank r, let p be a lexicographic extension of $\mathcal{M}$ on $[a_1^{s_1}, a_2^{s_2}, \dots, a_r^{s_r}]$, and let q be a lexicographic extension of $\mathcal{M} \cup p$ on $[p^-, a_1^-, \dots, a_{r-1}^-]$. If $\tilde{\mathcal{M}} = \mathcal{M} \cup p \cup q$, then $\operatorname{disc}(\tilde{\mathcal{M}}) = \operatorname{disc}(\mathcal{M})$.*

Proof. We prove that $\operatorname{disc}(\tilde{\mathcal{M}}) = \operatorname{disc}(\mathcal{M})$ by checking that every cocircuit $\tilde{C}$ of $\tilde{\mathcal{M}}$ has the same discrepancy as some cocircuit of $\mathcal{M}$.

If $\tilde{C}(p) \neq 0$ and $\tilde{C}(q) \neq 0$ then, by the definition of lexicographic extension, there is a cocircuit C of $\mathcal{M}$ such that $\tilde{C}|_{\mathcal{M}} = C$ and $\tilde{C}(p) = -\tilde{C}(q)$. Hence $|\tilde{C}^+| = |C^+| + 1$ and $|\tilde{C}^-| = |C^-| + 1$.

The cocircuits $\tilde{C}$ with $\tilde{C}(p) = 0$ correspond to cocircuits of $\tilde{\mathcal{M}}/p$, and those with $\tilde{C}(q) = 0$ correspond to cocircuits of $\tilde{\mathcal{M}}/q$. Proposition 2.6 tells us that

$$\tilde{\mathcal{M}}/p \simeq \tilde{\mathcal{M}}/q \simeq \mathcal{M}/a_1[a_2^{-s_1 s_2}, \dots, a_r^{-s_1 s_2}][p^-, a_2^-, \dots, a_{r-1}^-],$$

and the result follows by induction on r (it is trivial for $r = 1$). $\square$

In a vector configuration V, the proof that $V[p][q]$ has the same discrepancy as V is very easy to understand. For every hyperplane h spanned by V, if $p \in h^+$ then $q \in h^-$, and hence q balances the discrepancy created by p. For a hyperplane h that goes through p but neither a_1 nor q, the fact that p and a_1 are inseparable implies that except for a_1, h looks like a hyperplane spanned by V going through a_1. Hence q must balance the discrepancy created by a_1. For hyperplanes that go through p and a_1 but not a_2 nor q, q balances the discrepancy created by a_2. And so on.

And as a direct corollary:

Theorem 4.2. *Let $\mathcal{M}$ be a uniform balanced oriented matroid of rank r, let p be a lexicographic extension on $[a_1^{s_1}, a_2^{s_2}, \dots, a_r^{s_r}]$, and let q be a lexicographic extension of $\mathcal{M} \cup p$ on $[p^-, a_1^-, \dots, a_{r-1}^-]$. Then $\mathcal{M} \cup p \cup q$ is balanced.*

This provides the following construction to construct balanced matroids (and hence, by duality, to construct neighborly matroids). Let $\mathcal{M}_0 = \mathcal{D}_r$ be the oriented matroid associated to the totally cyclic vector configuration of rank r of minimal cardinality, which is balanced. And let

$$x_k = [e_{k1}^{s_{k1}}, \dots, e_{kr}^{s_{kr}}], \qquad y_k = [x_k^-, e_{k1}^-, \dots, e_{k(r-1)}^-], \qquad \mathcal{M}_k = \mathcal{M}_{k-1}[x_k][y_k],$$

where each e_{ij} is a different element of $\mathcal{M}_{i-1}$. Then $\mathcal{M}_k$ is balanced, and $\mathcal{M}_k^*$ neighborly. Since $\mathcal{M}_0$ is realizable, $\mathcal{M}_k^*$ is too, and hence $\mathcal{M}_k^*$ represents a neighborly polytope.

We call the double extension of Theorem 4.2 *Gale sewing*, and we denote by $\mathcal{G}$ the family of polytopes whose dual is constructed by repeatedly Gale sewing from some $\mathcal{D}_r$.

Corollary 4.3. *Let $\mathcal{P}$ be a neighborly matroid of rank d and n elements, then there exists a neighborly matroid $\tilde{\mathcal{P}}$ of rank $d+2$ with $n+2$ elements with two distinguished elements x and y such that $\tilde{\mathcal{P}}/\{x,y\} = \mathcal{P}$.*

Cyclic polytopes are a first example of polytopes in $\mathcal{G}$.

Proposition 4.4. *Let $\mathcal{M}$ be an oriented matroid dual to a cyclic polytope $c(d,n)$, with elements $s_1, s_2, \dots, s_n$ labeled according to the cyclic order. Then the extension $\mathcal{M} \cup \{x\} = \mathcal{M}[s_n^-, s_{n-1}^-, \dots, s_d^-]$ is an oriented matroid dual to the cyclic polytope $c(d+1, n+1)$, with $x = s_{n+1}$.*

Proof. The circuits of the cyclic polytope are as follows: the circuits X and Y supported by the $r+1$ points $e_1 < e_2 < \cdots < e_{r+1}$ are $X(e_i) = (-1)^i$ and $Y(e_i) = (-1)^{i+1}$ [6, Section 9.4].

If C is a cocircuit of $\mathcal{M}[x]$ (hence a circuit of its dual), the signature of the lexicographic extension implies that $C(x)$ is opposite to the sign of the largest non-zero element. And thus $\mathcal{M} \cup x$ is dual to $c(d+1, n+1)$. □

When Gale sewing, one can interchange the role of a_1, p and q as follows.

Lemma 4.5. *Let $\mathcal{M}$ be a uniform oriented matroid on a ground set E, and consider the lexicographic extensions*

$$p = [a_1^{s_1}, \dots, a_r^{s_r}], \qquad q = [p^-, a_1^-, \dots, a_{r-1}^-];$$
$$p' = [a_1^{-s_1}, \dots, a_r^{-s_r}], \qquad q' = [p'^-, a_1^-, \dots, a_{r-1}^-];$$
$$p'' = [a_1^+, a_2^{-s_1 s_2}, \dots, a_r^{-s_1 s_r}], \qquad q'' = [p''^-, a_1^-, \dots, a_{r-1}^-].$$

Then,

$$\mathcal{M}[p][q] \overset{\varphi}{\simeq} \mathcal{M}[p'][q'], \text{ and} \tag{4}$$

$$\mathcal{M}[p][q] \overset{\psi}{\simeq} \mathcal{M}[p''][q''], \tag{5}$$

where $\varphi(p) = q'$, $\varphi(q) = p'$ and $\varphi(e) = e$ for $e \in E$. If $s_1 = +$, then $\psi(p) = a_1$, $\psi(q) = q''$, $\psi(a_1) = p''$ and $\psi(e) = e$ for $e \in E \setminus \{a_1\}$, and if $s_1 = -$, then $\psi(p) = q''$, $\psi(q) = a_1$, $\psi(a_1) = p''$ and $\psi(e) = e$ for $e \in E \setminus \{a_1\}$

Proof. We start proving $\mathcal{M}[p][q] \overset{\varphi}{\simeq} \mathcal{M}[p'][q']$. That is, for every cocircuit $C \in \mathcal{C}^*(\mathcal{M}[p][q])$ we want to find a cocircuit $C' \in \mathcal{C}^*(\mathcal{M}[p'][q'])$ with $C'(\varphi(a)) = C(a)$ for all $a \in E \cup \{p, q\}$, that is $C'(p') = C(q)$, $C'(q') = C(p)$ and $C'(e) = C(e)$ for $e \in E$. Let D be the restiction of C to E.

If $C(q) \neq 0$ and $C(p) \neq 0$, let i be minimal with $D(a_i) \neq 0$. By construction, $C(p) = s_i D(a_i)$ and $C(q) = -C(p) = -s_i D(a_i)$. By the definition of $\mathcal{M}[p'][q']$, there is a cocircuit C' that expands D, with $C'(p') = -s_i D(a_i)$ and $C'(q') = -C(p') = s_i D(a_i)$.

To deal with the case when $C(q) = 0$ or $C(p) = 0$, by induction it is direct to see that $\mathcal{M}[p][q]/p \simeq \mathcal{M}[p'][q']/q'$ and that $\mathcal{M}[p][q]/q \simeq \mathcal{M}[p'][q']/p'$.

Now, to see that $\mathcal{M}[p][q] \overset{\psi}{\simeq} \mathcal{M}[p''][q'']$ we can assume that $s_1 = +$, otherwise we use eq. (4) to change the order of p and q. In this case we will prove that $\mathcal{M}[p] \simeq \mathcal{M}[p'']$. This suffices to prove eq. (5) since $[p^-, a_1^-, \dots, a_{r-1}^-]$ and $[a_1^-, p^-, \dots, a_{r-1}^-]$ give the same signature when p and a_1 are contravariant.

For every cocircuit $C \in \mathcal{C}^*(\mathcal{M}[p])$ we want to find now a cocircuit $C'' \in \mathcal{C}^*(\mathcal{M}[p''])$ with $C''(p'') = C(a_1)$, $C''(a_1) = C(p'')$ and $C''(e) = C(e)$ for $e \in E \setminus \{a_1\}$. Again, let D be the restriction of C to E.

If $C(p) \neq 0$ and $C(a_1) \neq 0$, then $C(p) = C(a_1) = D(a_1)$. And D can also be expanded to a cocircuit C'' of $\mathcal{M}[p'']$ with $C''(p'') = C''(a_1) = D(a_1)$. For circuits with $C(p) \neq 0$ and $C(a_1) = 0$, then $\mathcal{M}[p]/a_1 = \mathcal{M}[a_2^{s_2}, \dots, a_r^{s_r}] \simeq \mathcal{M}[p'']/p''$ by Proposition 2.6. Finally, if $C(p) = 0$ and $C(a_1) \neq 0$, then, again by Proposition 2.6, $\mathcal{M}[p]/p \simeq \mathcal{M}[a_2^{-s_2}, \dots, a_r^{-s_r}] = \mathcal{M}[p'']/a_1$. ☐

Using Proposition 2.6, we obtain the following result that explains the contractions of Gale sewing extensions:

Lemma 4.6. *Let $\mathcal{M}$ be a uniform oriented matroid of rank r, and consider the lexicographic extensions*

$$\begin{aligned} p &= [a_1^{s_1}, a_2^{s_2}, \dots, a_r^{s_r}], & q &= [p^-, a_1^-, \dots, a_{r-1}^-]; \\ p' &= [a_2^{-s_1 s_2}, \dots, a_r^{-s_1 s_r}], & q' &= [p'^-, \dots, a_{r-1}^-]. \end{aligned}$$

Then $(\mathcal{M}[p][q])/q \simeq (\mathcal{M}/a_1)[p'][q']$.

Proof. Repeatedly applying Proposition 2.6,

$$\begin{aligned} (\mathcal{M}[p][q])/q &= \big(\mathcal{M}\underbrace{[a_1^{s_1}, a_2^{s_2}, \dots, a_r^{s_r}]}_{p}\underbrace{[p^-, a_1^-, \dots, a_{r-1}^-]}_{q}\big)/q \\ &\overset{\varphi}{\simeq} \big(\mathcal{M}\underbrace{[a_1^{s_1}, a_2^{s_2}, \dots, a_r^{s_r}]}_{p}/p\big)\underbrace{[a_1^-, \dots, a_{r-1}^-]}_{\varphi(p)=q'} \\ &\overset{\psi}{\simeq} \big(\mathcal{M}/a_1\big)\underbrace{[a_2^{-s_1 s_2}, \dots, a_r^{-s_1 s_r}]}_{\psi(a_1)=p'}\underbrace{[\psi(a_1)^-, \dots, a_{r-1}^-]}_{q'}. \end{aligned}$$

☐

In fact, from this result and Proposition 2.6, it is easy to derive that subpolytopes of Gale sewn polytopes are also Gale sewn:

Proposition 4.7. *Let $\mathcal{P}$ be a neighborly polytope in $\mathcal{G}$, and let e be an element of $\mathcal{P}$. Then $\mathcal{P} \setminus e$ is also in $\mathcal{G}$.*

Proof. The proof is by induction on the rank of $\mathcal{P}$. When $\mathcal{P}$ has rank 0 then $\mathcal{P}^* = \mathcal{D}_r$ and $(\mathcal{P} \setminus e)^* = \mathcal{D}_r/e = \mathcal{D}_{r-1}$.

Otherwise, let $\mathcal{M} = \mathcal{P}^*$. That $\mathcal{P}$ belongs to $\mathcal{G}$ means that there is a matroid $\mathcal{N}$ whose dual $\mathcal{N}^*$ corresponds to a polytope $\mathcal{Q}$ in $\mathcal{G}$, such that $\mathcal{M} = \mathcal{N}[p][q]$ where $p = [a_1^{s_1}, a_2^{s_2}, \dots, a_r^{s_r}]$ and $q = [p^-, a_1^-, \dots, a_{r-1}^-]$. We will show that for every element e of $\mathcal{P}$, there is some element e' of $\mathcal{N}$ such that $(\mathcal{P} \setminus e)^* \simeq (\mathcal{N}/e')[p'][q'] = (\mathcal{Q} \setminus e')^*[p'][q']$. Since $\operatorname{rank} \mathcal{Q} = \operatorname{rank} \mathcal{P} - 2$, by the induction hypothesis $(\mathcal{Q} \setminus e') \in \mathcal{G}$ and the result follows.

If $e = q$, then by Lemma 4.6, $(\mathcal{P} \setminus e)^* = (\mathcal{N}[p][q])/q \simeq (\mathcal{N}/a_1)[p'][q']$. The case $e = p$ is analogous because of Lemma 4.5. If $e = a_i$, then $(\mathcal{P} \setminus e)^* \simeq (\mathcal{N}/a_i)[p'][q']$, where p' and q' have the same signature as p and q but omitting the element a_i. For the remaining elements e, $(\mathcal{P} \setminus e)^* \simeq (\mathcal{N}/e)[p][q]$. ☐

4.1. **Combinatorial description of the polytopes in $\mathcal{G}$.** Let P be a polytope that defines an acyclic oriented matroid $\mathcal{P}$, and let $\mathcal{M} = \mathcal{P}^*$ be its dual matroid. The essence of Gale sewing is to construct a new polytope $\tilde{P}$ whose matroid $\tilde{\mathcal{P}}$ is dual to $\tilde{\mathcal{M}} = \mathcal{M}[p]$, a lexicographic extension of $\mathcal{M}$ on $p = [a_1^{s_1}, a_2^{s_2}, \dots, a_k^{s_k}]$. In this section we will see that the combinatorics of $\tilde{P}$ are described by *lexicographic triangulations* of P.

Remark 4.8. The resemblance with the description of the vertex figures of the neighborly cubical polytopes in [23] is not a coincidence. Indeed, the duals of those vertex figures are lexicographic extensions of the dual of a fixed neighborly polytope.

Let $v_1, \dots, v_n$ be the vertices of $P \subset \mathbb{R}^d$, and assume (reordering the v_i's if necessary) that a_i is the dual element corresponding to v_i. Let V be the $n \times dn$ matrix whose rows list the coordinates of the v_i's:

$$V = \begin{bmatrix} \text{———} \; v_1 \; \text{———} \\ \text{———} \; v_2 \; \text{———} \\ \vdots \\ \text{———} \; v_n \; \text{———} \end{bmatrix}.$$

Then the following $(n+1) \times (d+1)$ matrix $\tilde{V}$ encodes the vertices of $\tilde{P}$, for some small enough ε.

$$\tilde{V} = \left[\begin{array}{c|c} \text{———} \; v_1 \; \text{———} & -s_1 \\ \text{———} \; v_2 \; \text{———} & -s_2\varepsilon \\ \text{———} \; v_3 \; \text{———} & -s_3\varepsilon^2 \\ \vdots & \vdots \\ \text{———} \; v_k \; \text{———} & -s_k\varepsilon^{k-1} \\ \text{———} \; v_{k+1} \; \text{———} & 0 \\ \vdots & \vdots \\ \text{———} \; v_n \; \text{———} & 0 \\ \text{———} \; \mathbf{0} \; \text{———} & 1 \end{array}\right].$$

The vertex figure of p in $\tilde{P}$ is combinatorially equivalent to P, hence the faces of $\tilde{P}$ that contain p are isomorphic to pyramids over faces of P. On the other hand, the faces of $\tilde{P}$ that do not contain p correspond to faces of a regular subdivision of P: a *lexicographic subdivision* of P on $[v_1^{-s_1}, v_2^{-s_2}, \dots, v_k^{-s_k}]$. When $k = r$, then this subdivision is a triangulation.

Our formulation of the definition of lexicographic subdivision is based on [11]. However we use a different ordering, the same as in [22], that mirrors the definition of lexicographic extension. See also [18].

Definition 4.9. Let P be a d-polytope with n vertices $v_1, \dots, v_n$. The *lexicographic subdivision* of P on $[v_1^{s_1}, v_2^{s_2}, \dots, v_k^{s_k}]$, where $s_i = \pm 1$, is defined recursively as follows.

- If $s_1 = +1$ (pushing), then the lexicographic subdivision of P is the union of the lexicographic subdivision of $P \setminus v_1$ on $[v_2^{s_2}, \dots, v_k^{s_k}]$, and the simplices joining v_1 to the (lexicographically subdivided) faces of $P \setminus v_1$ visible from it.
- If $s_1 = -1$ (pulling), then the lexicographic subdivision of P is the unique subdivision in which every maximal cell contains v_1 and which, restricted to each proper face F of P, coincides with the lexicographic subdivision of that face.



FIGURE 2. The lifting of a pentagon P to $\tilde{P}$ when $p = [a_1^-, a_4^+]$. Its upper envelope are pyramids over facets of P, while the lower envelope is the lexicographic triangulation of P on $[a_1^+, a_4^-]$.

5. COMPARING AND COMBINING THE CONSTRUCTIONS

In this section we compare and combine the two constructions, which are strongly related. The first goal is to prove that all neighborly polytopes in $\mathcal{O}$ belong to $\mathcal{G}$. From Proposition 4.7 one deduces that to prove that $\mathcal{O} \subseteq \mathcal{G}$ it is enough to see that $\mathcal{E} \subseteq \mathcal{G}$. As a first step we will prove that if we sew on a neighborly polytope in $\mathcal{G}$ (through a specific universal flag), we obtain a new polytope in $\mathcal{G}$.

Let $\mathcal{P}$ be a neighborly matroid in $\mathcal{G}$. That means that its dual matroid $\mathcal{M} = \mathcal{P}^*$ is constructed using Gale sewing as in Section 4. Specifically, $\mathcal{M} = \mathcal{M}_m$ where $\mathcal{M}_0 = \mathcal{D}_r$ is a minimal totally cyclic configuration $\{a_0, \dots, a_r\}$ of rank r and

$$\mathcal{M}_k = \mathcal{M}_{k-1}[x_k][y_k],$$

with x_k and y_k being the lexicographic extensions

$$x_k = [e_{k1}^{s_{k1}}, \dots, e_{kr}^{s_{kr}}], \qquad\qquad y_k = [x_k^-, e_{k1}^-, \dots, e_{k(r-1)}^-].$$

Let $T_j = \bigcup_{i=0}^{j-1}\{x_{m-i}, y_{m-i}\}$, that is $T_{m-k} = \{x_m, y_m, \dots, x_{k+1}, y_{k+1}\}$. And let $\mathcal{T} = \{T_i\}_{i=1}^m$. Define $\mathcal{P}_k = \mathcal{M}_k^*$, and observe that $\mathcal{P}_k = \mathcal{P}/T_{m-k}$ for $k = 0, \dots, m$. By construction, $\mathcal{T}$ is a universal flag of $\mathcal{P}$. Let $\mathcal{T}'$ be some flag that contains $\mathcal{T}$. Thanks to Lemma 4.5 we can assume without loss of generality that all split faces in $\mathcal{T}'$ are y-split.

We can apply the extended sewing construction of Theorem 3.7 to $\mathcal{P}$ and $\mathcal{T}'$. Let $\tilde{\mathcal{P}}$ be $\mathcal{P} \cup p$, with p sewn onto $\mathcal{P}$ through $\mathcal{T}'$. We define $\tilde{T}_{j+1} = T_j \cup y_{m-j} \cup p$, that is $\tilde{T}_{m-k} = \{x_m, y_m, \dots, x_{k+2}, y_{k+2}\} \cup \{y_{k+1}\} \cup \{p\}$. Then $\tilde{\mathcal{T}} = \{\tilde{T}_i\}_{i=1}^m$ is a universal flag of $\tilde{\mathcal{P}}$ by Proposition 3.9. We denote $\tilde{\mathcal{P}}_k = \tilde{\mathcal{P}}/\tilde{T}_{m-k}$ and observe that $\tilde{\mathcal{P}}_k \simeq \mathcal{P}_k[\mathcal{T}'/T_{m-k}]$ where the sewn vertex is x_{k+1}, and thus $\tilde{\mathcal{P}}_k \backslash x_{k+1} \simeq \mathcal{P}_k$. To provide shorter proofs, we will sometimes refer to p as x_{m+1}.

Finally, let $\tilde{\mathcal{M}} = \tilde{\mathcal{M}}_m$ be the oriented matroid constructed as follows:

$$\begin{aligned}
\tilde{\mathcal{M}}_0 &= \mathcal{D}_{r+1} \text{ on a ground set } \{\tilde a_1, \dots, \tilde a_{r+1}, \tilde x_1\},\\
\tilde y_k &= \begin{cases} [\tilde x_k^+, (\tilde e_{k1})^{-s_{k1}}, \dots, (\tilde e_{kr})^{-s_{kr}}] & \text{if } T_{m-k+1} \text{ is not split in } \mathcal{T}',\\ [\tilde x_k^-, (\tilde e_{k1})^{s_{k1}}, \dots, (\tilde e_{kr})^{s_{kr}}] & \text{if } T_{m-k+1} \text{ is split in } \mathcal{T}'.\end{cases}\\
\tilde x_{k+1} &= [\tilde y_k^-, (\tilde e_{k1})^-, \dots, (\tilde e_{k(r-1)})^-],\\
\tilde{\mathcal{M}}_k &= \tilde{\mathcal{M}}_{k-1}[\tilde y_k][\tilde x_{k+1}],
\end{aligned}$$

where if $e_{ij} = x$ then $\tilde e_{ij}$ means the element labeled as $\tilde x$.

Then we claim that $\tilde{\mathcal{M}}$ is the dual configuration of $\tilde{\mathcal{P}}$, and hence $\tilde{\mathcal{P}}$ is in $\mathcal{G}$.



FIGURE 3. Example: $\mathcal{M}$ is constructed from $\{A, B\}$ after Gale sewing $C = [B^+]$ and $D = [C^-]$. In its dual $\mathcal{P} = \mathcal{M}^*$, $\{C, D\}$ is a universal edge. $\tilde{\mathcal{P}} = \mathcal{P}[E]$ is obtained by sewing E on $\mathcal{P}$ through $\{C, D\}$. Its dual matroid, $\tilde{\mathcal{M}} = \tilde{\mathcal{P}}^*$ can be constructed from $\{A, B, C\}$ and Gale sewing $D = [C^+, B^-]$ and $E = [D^-, C^-]$.

Proposition 5.1. *With the notations from above, $\tilde{\mathcal{M}} \simeq \tilde{\mathcal{P}}^*$ via the isomorphism $x \mapsto \tilde x$.*

Proof. We will prove that $\tilde{\mathcal{M}}_k \simeq \tilde{\mathcal{P}}_k^*$, for all k. The proof will use induction on k and assume that $\tilde{\mathcal{M}}_{k-1} \simeq \tilde{\mathcal{P}}_{k-1}^*$. It is straightforward that $\tilde{\mathcal{M}}_0 \simeq \tilde{\mathcal{P}}_0^*$ since $\mathcal{P}_0$ is 0-dimensional.

We will use Theorem 2.1 twice. Specifically, we will use that if $\mathcal{M} \setminus p \simeq \mathcal{N} \setminus p$ and $\mathcal{M}/p \simeq \mathcal{N}/p$, then $\mathcal{M}$ and $\mathcal{N}$ are the same oriented matroid up to the reorientation of p. If additionally there is an element q such that p and q are covariant (or contravariant) in $\mathcal{N}$ and $\mathcal{M}$, then $\mathcal{M} \simeq \mathcal{N}$.

In particular, we will prove that $\tilde{\mathcal{M}}_k/\tilde x_{k+1} \simeq \tilde{\mathcal{P}}_k^*/x_{k+1}$ and that $\tilde{\mathcal{M}}_k \setminus \tilde x_{k+1} \simeq \tilde{\mathcal{P}}_k^* \setminus x_{k+1}$. Then the claim $\tilde{\mathcal{M}} \simeq \tilde{\mathcal{P}}^*$ will follow directly from the fact that $\tilde x_{k+1}$ and $\tilde y_k$ are covariant in $\tilde{\mathcal{M}}_k$ and that x_{k+1} and y_k are covariant in $\tilde{\mathcal{P}}_k^*$.

$\underline{\tilde{\mathcal{M}}_k/\tilde x_{k+1} \simeq \tilde{\mathcal{P}}_k^*/x_{k+1}}$

Since $\tilde{\mathcal{P}}_k^*/x_{k+1} \simeq \mathcal{P}_k^* = \mathcal{M}_k$ because $\tilde{\mathcal{P}}_k \setminus x_{k+1} \simeq \mathcal{P}_k$, we just need to prove that

$$\tilde{\mathcal{M}}_k/\tilde{x}_{k+1} \simeq \mathcal{M}_k. \tag{6}$$

By Lemma 4.6, $(\tilde{\mathcal{M}}_k/\tilde{x}_{k+1}) \simeq (\tilde{\mathcal{M}}_{k-1}/\tilde{x}_k)[x'_k][y'_k]$, where $x'_k = [\tilde{e}_{k1}^{s_{k1}},\dots,\tilde{e}_{kr}^{s_{kr}}]$ and $y'_k = [{x'_k}^-, \tilde{e}_{k1}^-,\dots,\tilde{e}_{k(r-1)}^-]$. Using that $\tilde{\mathcal{M}}_0/\tilde{x}_1 \simeq \mathcal{M}_0$ we get the desired result by induction on k.

$\tilde{\mathcal{M}}_k \setminus \tilde{x}_{k+1} \simeq \tilde{\mathcal{P}}_k^* \setminus x_{k+1}$

The first step is to prove that $(\tilde{\mathcal{M}}_k \setminus \tilde{x}_{k+1}) \setminus \tilde{y}_k \simeq (\tilde{\mathcal{P}}_k^* \setminus x_{k+1}) \setminus y_k$. Indeed, using the induction hypothesis,

$$(\tilde{\mathcal{M}}_k \setminus \tilde{x}_{k+1}) \setminus \tilde{y}_k = \tilde{\mathcal{M}}_{k-1} \simeq \tilde{\mathcal{P}}_{k-1}^* = (\tilde{\mathcal{P}}_k/\{y_k, x_{k+1}\})^* = (\tilde{\mathcal{P}}_k^* \setminus x_{k+1}) \setminus y_k.$$

Now we prove that $(\tilde{\mathcal{M}}_k \setminus \tilde{x}_{k+1})/\tilde{y}_k \simeq (\tilde{\mathcal{P}}_k^* \setminus x_{k+1})/y_k$. First, using Lemma 2.5 and (6), we see that

$$(\tilde{\mathcal{M}}_k \setminus \tilde{x}_{k+1})/\tilde{y}_k \simeq (\tilde{\mathcal{M}}_k/\tilde{x}_{k+1}) \setminus \tilde{y}_k \simeq \mathcal{M}_k \setminus y_k.$$

Now, using again Lemma 2.5 and that $\tilde{\mathcal{P}}_k \setminus x_{k+1} \simeq \mathcal{P}_k$, we see that

$$(\tilde{\mathcal{P}}_k^* \setminus x_{k+1})/y_k = (\tilde{\mathcal{P}}_k/x_{k+1} \setminus y_k)^* \simeq (\tilde{\mathcal{P}}_k \setminus x_{k+1}/y_k)^* = (\mathcal{P}_k/y_k)^* = {\mathcal{P}_k}^* \setminus y_k.$$

Our claim follows since $\mathcal{M}_k = \mathcal{P}_k^*$ by definition.

Because of Theorem 2.1, so far we have seen that $\tilde{\mathcal{M}}_k \setminus \tilde{x}_{k+1} \simeq \tilde{\mathcal{P}}_k^* \setminus x_{k+1}$ up to reorientation of y_k. We will conclude the proof by seeing that y_k and $\tilde{y}_k$ have the same orientation in $\mathcal{M}_k$ and in $\tilde{\mathcal{M}}_k$ respectively.

If T_{m-k+1} is not split, then $\tilde{y}_k$ is contravariant with $\tilde{x}_k$ in $\tilde{\mathcal{M}}_k$ by construction. Moreover, y_k is contravariant with x_k in $(\tilde{\mathcal{P}}_k/x_{k+1})^*$ since they are covariant in the primal: by Proposition 2.6

$$\tilde{\mathcal{P}}_k/x_{k+1} \simeq \big(\mathcal{P}_k \underbrace{[\mathcal{T}'/T_{m-k}]}_{x_{k+1}}\big)/x_{k+1} \simeq \big(\mathcal{P}_k/x_k\big)\underbrace{[y_k^-,\dots]}_{x_k},$$

where the last isomorphism sends x_k to the sewn vertex, which is $(+1)$-inseparable from y_k.

If T_{m-k+1} is split, then $\tilde{y}_k$ is covariant with $\tilde{x}_k$ in $\tilde{\mathcal{M}}_k$. Moreover, y_k is covariant with x_k in $(\tilde{\mathcal{P}}_k/x_{k+1})^*$ because $\tilde{\mathcal{P}}_k/x_{k+1} \simeq \mathcal{P}_k/x_k[y_k^+,\dots]$.

□

There is a missing detail to conclude that $\mathcal{E} \subseteq \mathcal{G}$ from Proposition 5.1: we have to check that after sewing, the universal flags of Proposition 3.9 are also of the form $T_j = \cup_{i=0}^{j-1}(x_{m-i}, y_{m-i})$ for some Gale sewn configuration.

This is consequence of Lemma 4.5, which allows to change the order of the sewings in $\tilde{\mathcal{M}}$.

Remark 5.2. The fact that $\mathcal{E} \subsetneq \mathcal{G}$ implies that in some sense Gale sewing generalizes ordinary sewing. However, it is not true that Theorem 3.5 is a consequence of Theorem 4.2, because there are neighborly matroids that have universal flags but are not in $\mathcal{G}$. Hence one can sew on them but they cannot be treated with Proposition 5.1. This will become clear in Section 5.2, where both constructions are needed.

5.1. Some exact numbers.

Fix n and $d = 2m$. We have worked with four families of neighborly d-polytopes with n vertices:

- **$\mathcal{N}$:** All neighborly d-polytopes with n vertices.
- **$\mathcal{G}$:** All Gale sewn d-polytopes with n vertices.
- **$\mathcal{S}$:** All totally sewn d-polytopes with n vertices.
- **$\mathcal{E}$:** All d-polytopes with n vertices obtained with the extended sewing construction.
- **$\mathcal{O}$:** All d-polytopes with n vertices obtained by extended sewing and omitting.

Table 1 contains the exact number of combinatorial types in each of these families for the cases $d = 4$ and $n = 8, 9$ and for $d = 6$ and $n = 10$. Exact numbers for $\mathcal{N}$ come from [3] and [8], exact numbers for $\mathcal{S}$ and $\mathcal{O}$ come from [24]. Numbers for $\mathcal{G}$ and $\mathcal{E}$ have been computed with the help of `polymake` [13].

TABLE 1. Exact number of combinatorial types

d	n	$\mathcal{S}$	$\mathcal{E}$	$\mathcal{O}$	$\mathcal{G}$	$\mathcal{N}$
4	8	3	3	3	3	3
4	9	18	18	18	18	23
6	10	15	26	28	28	37

In view of Table 1, the relationships that we know between these families are summarized in the following proposition.

Proposition 5.3. $\mathcal{S} \subsetneq \mathcal{E} \subsetneq \mathcal{O} \subseteq \mathcal{G} \subsetneq \mathcal{N}$.

Which yields the following question.

Question 1. *Is $\mathcal{O} = \mathcal{G}$?*

5.2. Non-realizable neighborly oriented matroids.

Because the only neighborly oriented matroids of rank 3 are cyclic polytopes, there are no non-realizable neighborly oriented matroids of rank 3. However, the sphere "M^{10}_{425}" from Altshuler's list [2] corresponds to a neighborly matroid of rank 5 with 10 elements. In [7], this matroid is proved to be non-realizable, giving a proof that there exist non-realizable neighborly matroids. Kortenkamp's construction [16] can be used to build non-realizable neighborly matroids of corank 3. We show that there are many non-realizable neighborly matroids.

Theorem 5.4. *There exists a non-realizable neighborly matroid of rank d with n elements for every $d \geq 5$ and $n \geq d + 5$.*

Proof. We start with M^{10}_{425}. With the vertex labeling of [2], $\{0, 1\}$, $\{2, 3\}$, $\{4, 5\}$, $\{6, 7\}$ and $\{8, 9\}$ are universal edges of M^{10}_{425} because the corresponding contractions are polygons with 8 vertices. In particular, $\{0, 1\} \subset \{0, 1, 2, 3\}$ is a universal flag. Hence, applying Theorem 3.7 we get many non-realizable matroids of rank 5 with n vertices for any $n \geq 10$.

Now, applying Corollary 4.3 to these matroids, we get non-realizable oriented matroids of rank $5 + 2k$ and n vertices for any $k \geq 0$ and any $n \geq 10 + 2k$.

To get non-realizable matroids of even rank, just observe that any single element extension on the dual of a neighborly matroid of rank $2k + 1$ yields the dual of a neighborly matroid of rank $2k + 2$. □

All neighborly matroids of rank $2m+1$ that have $n \leq 2m+3$ vertices are cyclic polytopes. Moreover, all oriented matroids of rank 5 with 8 elements are realizable [6, Corollary 8.3.3]. Hence the first case that Theorem 5.4 does not deal with are neighborly matroids of rank 5 with 9 elements.

Question 2. *Do there exist non-realizable neighborly oriented matroids of rank* 5 *on* 9 *elements?*

6. Bounds on the number of neighborly polytopes

The aim of this section is to find lower bounds for $\mathrm{nb}_l(n,d)$, the number of combinatorial types of vertex-labeled neighborly polytopes with n vertices in dimension d. Because of Theorem 2.8, two neighborly polytopes with the same combinatorial type have the same oriented matroid. Therefore, it is enough to focus on giving bounds for the numbers of labeled realizable neighborly matroids. In the remainder of this paper, all polytopes and oriented matroids will be labeled. Our bounds will be so large that the naive bounds for unlabeled combinatorial types obtained by dividing by $n!$ would have similar asymptotics.

More precisely, we will use the Gale sewing technique of Theorem 4.2 to give lower bounds on the number of d-polytopes with n vertices that are in $\mathcal{G}$.

A first step is to compute lower bounds for $\ell_l(n,r)$, the smallest number of different labeled lexicographic extensions that a balanced matroid of rank r with n elements can have. There are $2^r\frac{n!}{(n-r)!}$ different expressions for lexicographic extensions of a matroid $\mathcal{M}$ of rank r and n elements, yet not all of them represent different labeled oriented matroids. We aim to avoid counting the same extension twice with two different expressions.

Proposition 6.1. *Let $\mathcal{M}$ be a rank r labeled balanced matroid with $n = r+1+2m$ elements. The number of different labeled lexicographic extensions of $\mathcal{M}$ fulfills*

$$\ell_l(n,r) \geq 2 \cdot \frac{n!}{(n-r+1)!}. \tag{7}$$

Proof. We focus only on those extensions where $s_i = +$ for all i. Those are unambiguous except for the last element $a_r^{s_r}$.

For this, observe that if $[a_1^+,\dots,a_r^+]$ and $[{a'_1}^+,\dots,{a'_r}^+]$, with $r>1$, provide the same oriented matroid, then a_1 and a'_1 must be contravariant. But balanced matroids of rank $r \geq 2$ and $r+1+2m$ elements only have covariant pairs (see Lemma 2.11). If a_i and a'_i are the first different elements and $i<r$, we can apply the previous argument on the contraction by $\{a_1,\dots,a_{i-1}\}$.

For the last element, observe that $\mathcal{M}/\{a_1,\dots,a_{r-1}\}$ is a matroid of rank 1, and that there are exactly two possible different extensions for a matroid of rank 1. □

Remark 6.2. In the bound of (7), we lose a factor of up to 2^{r-1} from the real number. This factor is asymptotically much smaller than $\frac{2n!}{(n-r+1)!}$. In fact, it is not difficult to prove that $\ell_l(n,r) \geq 2^{r-1}\frac{n!}{(n-1)(n-r)!}$ by giving some cyclic order to the elements of $\mathcal{M}$ and counting only the lexicographic extensions $[a_1^{s_1},\dots,a_r^{s_r}]$ that fulfill

(i) For $1<i<r$, $a_i^{s_i}$ is not $b^{-\alpha s_{i-1}}$ if $b<a_{i-1}$ and b and a_{i-1} are α-inseparable in $\mathcal{M}/\{a_1,\dots,a_{i-2}\}$.

(ii) For $1<i<r$, $a_i^{s_i}$ is not $c^{\alpha\beta s_{i-1}}$ when there exists b with $c<b<a_{i-1}$ such that b and a_{i-1} are α-inseparable in $\mathcal{M}/\{a_1,\dots,a_{i-2}\}$ and c is β-inseparable from b in $\mathcal{M}/\{a_1,\dots,a_{i-1}\}$.

(iii) a_r and a_{r-1} are α-inseparable in $\mathcal{M}/\{a_1, \dots, a_{r-2}\}$, $a_{r-1} < a_r$ and $s_r = \alpha s_{r-1}$.

But then the formulas become more complicated and and adds nothing substantial to the result.

Once we have bounds for $\ell_l(n, r)$, we can obtain bounds for $\mathrm{nb}_l(n, d)$. But first we do a case where we know the exact number.

Lemma 6.3. *The number of labeled balanced matroids of rank r with $r + 3$ elements is $\frac{1}{2}(r + 2)!$.*

Proof. Balanced matroids of rank r with $r+3$ elements are dual to polygons with $r+3$ vertices in $\mathbb{R}^2$. There are clearly $\frac{1}{2}(r + 2)!$ of them. $\square$

We will need the following result on the inseparability graph of oriented matroids. The *inseparability graph* of an oriented matroid $\mathcal{M}$, $IG(\mathcal{M})$, is the graph that has the elements of $\mathcal{M}$ as vertices and whose edges are the pairs of inseparable elements.

Theorem 6.4 ([9, Theorem 1.1])**.** *Let $\mathcal{M}$ be a rank r uniform oriented matroid with n elements.*

- *If $r \le 1$ or $r \ge n - 1$, then $IG(\mathcal{M})$ is the complete graph.*
- *If $r = 2$ or $r = n - 2$, then $IG(\mathcal{M})$ is an n-cycle.*
- *If $2 < r < n - 2$, then $IG(\mathcal{M})$ is either a n-cycle, or a disjoint union of chains.*

Lemma 6.5. *The number of labeled balanced matroids of rank r with $r + 1 + 2m$ elements fulfills*

$$\mathrm{nb}_l(2m + r + 1, 2m) \ge \mathrm{nb}_l(2m + r - 1, 2m - 2)\frac{r + 2m}{2}\ell_l(r + 2m - 1, r).$$

Proof. This bound is a consequence of Theorem 4.2. Fix a labeled balanced matroid $\mathcal{M}$ of rank r with $r + 1 + 2(m - 1)$ elements. There are at least $\ell_l(r + 2m - 1, r)$ different labeled lexicographic extensions on $\mathcal{M}$. Let $p = [a_1^{s_1}, \dots, a_r^{s_r}]$ be any such extension, and let q be the extension $q = [p^-, a_1^-, \dots, a_{r-1}^-]$. We count the ways of labeling $\mathcal{M}[p][q]$ such that p gets label 0, and the labeling of $\mathcal{M}[p][q]$ on $\mathcal{M}$ preserves the relative order of the original labeling of $\mathcal{M}$. There are clearly $r + 2m$ ways to do it, which correspond to the possible labels that can be given to q. Since p is inseparable from at most two elements (see Theorem 6.4), and p and q are inseparable, we have counted each of these labeled matroids at most twice. Which gives $\mathrm{nb}_l(2m + r + 1, 2m) \ge \mathrm{nb}_l(2m + r - 1, 2m - 2)\frac{r+2m}{2}\ell_l(r + 2m - 1, r)$. $\square$

Corollary 6.6. *The number of labeled balanced matroids of rank r with $r + 1 + 2m$ elements fulfills*

$$\mathrm{nb}_l(2m + r + 1, 2m) \ge \prod_{i=0}^{m-1} \frac{r + 2 + 2i}{2}\ell_l(r + 1 + 2i, r).$$

Combining Proposition 6.1 and Corollary 6.6, we get the following bound on the number of labeled balanced matroids of rank r with $r + 1 + 2m$ elements:

$$\mathrm{nb}_l(r + 1 + 2m, 2m) \ge \prod_{i=1}^{m} \frac{(r + 2i)!}{(2i)!}. \tag{8}$$

Theorem 6.7. *The number of labeled neighborly polytopes in dimension d with n vertices fulfills*

$$\mathrm{nb}_l(r+d+1,d) \geq \frac{(r+d)^{\left(\frac{r}{2}+\frac{d}{2}\right)^2}}{r^{\left(\frac{r}{2}\right)^2} d^{\left(\frac{d}{2}\right)^2} \mathrm{e}^{3\frac{r}{2}\frac{d}{2}}},$$

that is,

$$\mathrm{nb}_l(n,d) \geq \frac{(n-1)^{\left(\frac{n-1}{2}\right)^2}}{(n-d-1)^{\left(\frac{n-d-1}{2}\right)^2} d^{\left(\frac{d}{2}\right)^2} \mathrm{e}^{\frac{3d(n-d-1)}{4}}}.$$

Proof. First we assume that $d = 2m$ is even. We start from Equation (8), and try to approximate the natural logarithm of $\mathrm{nb}_l(r+1+2m,2m)$. Using that for any increasing function f, $\int_{s=a-1}^{b} f(s)\,\mathrm{d}s \leq \sum_{i=a}^{b} f(i)$, we get

$$\begin{aligned}
\ln\left(\mathrm{nb}_l(r+1+2m,2m)\right) &\geq \ln\left(\prod_{i=1}^{m}(r+2i)^{\underline{r}}\right) = \sum_{i=1}^{m}\sum_{j=0}^{r-1}\ln\left(r+2i-j\right)\\
&= \sum_{i=1}^{m}\sum_{j=1}^{r}\ln\left(2i+j\right) \geq \int_{i=0}^{m}\int_{j=0}^{r}\ln\left(2i+j\right)\mathrm{d}j\mathrm{d}i\\
&= \frac{(2m+r)^2\ln\left(2m+r\right)-r^2\ln\left(r\right)}{4} - m^2\ln\left(2m\right) - \frac{3mr}{2}.
\end{aligned}$$

Hence,

$$\mathrm{nb}_l(r+1+2m,2m) \geq \frac{(2\,m+r)^{\frac{1}{4}(2\,m+r)^2}}{r^{\frac{1}{4}r^2}\,(2\,m)^{m^2}\,\mathrm{e}^{\frac{3}{2}mr}}.$$

And therefore,

$$\mathrm{nb}_l(r+d+1,d) \geq \frac{(r+d)^{\left(\frac{r}{2}+\frac{d}{2}\right)^2}}{r^{\left(\frac{r}{2}\right)^2} d^{\left(\frac{d}{2}\right)^2} \mathrm{e}^{3\frac{r}{2}\frac{d}{2}}}. \tag{9}$$

If $d = 2m+1$, observe that any lexicographic extension of a balanced matroid with $r+1+2m$ elements in rank r will be balanced. Moreover, if this extension is of the form $p = [a_1^+, \dots]$ (like the extensions in Proposition 6.1), then p and a_1 are the only pair of contravariant elements in $\mathcal{M}$. Hence we can assign any of the possible $r+2m+2$ labels to p and we will have counted the extension at most twice. Therefore,

$$\begin{aligned}
\mathrm{nb}_l(r+2+2m,2m+1) &\geq \mathrm{nb}_l(r+1+2m,2m)\frac{2r+2m+2}{2}\ell_l(r+1+2m,r)\\
&\geq \mathrm{nb}_l(r+1+2m,2m)\prod_{j=0}^{r}(2+2m+j)
\end{aligned}$$

Now,

$$\begin{aligned}
\ln(\mathrm{nb}_l(r+2m+2,2m+1)) &\geq \int_{j=0}^{r}\int_{i=0}^{m}\ln\left(2i+j\right)\mathrm{d}i\mathrm{d}j + \int_{j=0}^{r}\ln\left(2m+2+j\right)\mathrm{d}j\\
&\geq \int_{j=0}^{r}\int_{i=0}^{m+\frac{1}{2}}\ln\left(2i+j\right)\mathrm{d}i\mathrm{d}j.
\end{aligned}$$

This last inequality is due to $\ln(2m+2+j) \geq \int_{i=m}^{m+\frac{1}{2}} \ln(2i+j)\,\mathrm{d}i$ and proves that Equation (9) also holds for odd dimension. □

Corollary 6.8. *If* $n > 2d$*, the number of labeled neighborly polytopes in dimension* d *with* n *vertices fulfills*

$$\mathrm{nb}_l(n,d) \geq \left(\frac{n-1}{\sqrt{\mathrm{e}^3}}\right)^{\frac{d(n-1)}{2}}.$$

Acknowledgements

I would like to thank Uli Wagner for stimulating discussions. Moreover, I am greatly indebted to Julian Pfeifle for his comments and corrections that greatly helped to improve the presentation.

References

[1] Noga Alon. The number of polytopes, configurations and real matroids. *Mathematika*, 33(1):62–71, 1986.

[2] Amos Altshuler. Neighborly 4-polytopes and neighborly combinatorial 3-manifolds with ten vertices. *Can. J. Math.*, 29:225–420, 1977.

[3] Amos Altshuler and Leon Steinberg. Neighborly 4-polytopes with 9 vertices. *J. Comb. Theory, Ser. A*, 15:270–287, 1973.

[4] David Barnette. A family of neighborly polytopes. *Isr. J. Math.*, 39:127–140, 1981.

[5] Tibor Bisztriczky. On sewing neighbourly polytopes. *Note Mat.*, 20(1):73–80, 2000/01.

[6] Anders Björner, Michel Las Vergnas, Bernd Sturmfels, Neil White, and Günter M. Ziegler. *Oriented matroids.* Encyclopedia of Mathematics and Its Applications. 46. Cambridge: Cambridge University Press. 516 p. , 1993.

[7] Jürgen Bokowski and Klaus Garms. Altshuler's sphere M^{10}_{425} is not polytopal. *Eur. J. Comb.*, 8:227–229, 1987.

[8] Jürgen Bokowski and Ido Shemer. Neighborly 6-polytopes with 10 vertices. *Isr. J. Math.*, 58:103–124, 1987.

[9] Raul Cordovil and Pierre Duchet. On sign-invariance graphs of uniform oriented matroids. *Discrete Math.*, 79(3):251–257, 1990.

[10] Raul Cordovil and Pierre Duchet. Cyclic polytopes and oriented matroids. *Eur. J. Comb.*, 21(1):49–64, 2000.

[11] Jesús A. De Loera, Jörg Rambau, and Francisco Santos. *Triangulations*, volume 25 of *Algorithms and Computation in Mathematics.* Springer-Verlag, Berlin, 2010. Structures for algorithms and applications.

[12] Lukas Finschi. *A Graph Theoretical Approach for Reconstruction and Generation of Oriented Matroids.* PhD thesis, ETH Zürich, 2001.

[13] Ewgenij Gawrilow and Michael Joswig. polymake: a framework for analyzing convex polytopes. In Gil Kalai and Günter M. Ziegler, editors, *Polytopes — Combinatorics and Computation*, pages 43–74. Birkhäuser, 2000.

[14] Jacob E. Goodman and Richard Pollack. Upper bounds for configurations and polytopes in $\mathbf{R}^d$. *Discrete Comput. Geom.*, 1:219–227, 1986.

[15] Branko Grünbaum. *Convex polytopes*, volume 221 of *Graduate Texts in Mathematics.* Springer-Verlag, New York, second edition, 2003. Prepared and with a preface by Volker Kaibel, Victor Klee and Günter M. Ziegler.

[16] Ulrich H. Kortenkamp. Every simplicial polytope with at most $d+4$ vertices is a quotient of a neighborly polytope. *Discrete Comput. Geom.*, 18(4):455–462, 1997.

[17] Michel Las Vergnas. Extensions ponctuelles d'une géométrie combinatoire orientee. Problèmes combinatoires et théorie des graphes, Orsay 1976, Colloq. int. CNRS No.260, 265-270 (1978)., 1978.

[18] Carl W. Lee. Regular triangulations of convex polytopes. In *Applied geometry and discrete mathematics*, volume 4 of *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, pages 443–456. Amer. Math. Soc., Providence, RI, 1991.

[19] Carl W. Lee and Matthew M. Menzel. A generalized sewing construction for polytopes. *Isr. J. Math.*, 176:241–267, 2010.
[20] Peter McMullen. The maximum numbers of faces of a convex polytope. *Mathematika, Lond.*, 17:179–184, 1970.
[21] Jürgen Richter-Gebert and Günter M. Ziegler. Zonotopal tilings and the Bohne-Dress theorem. In *Jerusalem combinatorics '93*, volume 178 of *Contemp. Math.*, pages 211–232. Amer. Math. Soc., 1994.
[22] Francisco Santos. Triangulations of oriented matroids. *Mem. Am. Math. Soc.*, 741:80 p., 2002.
[23] Raman Sanyal and Günter M. Ziegler. Construction and analysis of projected deformed products. *Discrete Comput. Geom.*, 43(2):412–435, 2010.
[24] Ido Shemer. Neighborly polytopes. *Isr. J. Math.*, 43:291–314, 1982.
[25] Bernd Sturmfels. Neighborly polytopes and oriented matroids. *Eur. J. Comb.*, 9(6):537–546, 1988.
[26] Ryan Trelford and Viktor Vigh. How to sew in practice? `arXiv:1102.4862v1`, 2011.

DEPT. MATEMÀTICA APLICADA II, UNIVERSITAT POLITÈCNICA DE CATALUNYA (UPC), BARCELONA, SPAIN.

E-mail address: `arnau.padrol@upc.edu`